

A GENERALIZATION OF THE SIMION-SCHMIDT BIJECTION FOR RESTRICTED PERMUTATIONS

Astrid Reifegerste

Institut für Mathematik, Universität Hannover

Welfengarten 1, D-30167 Hannover, Germany

reifegerste@math.uni-hannover.de

ABSTRACT. We consider the two permutation statistics which count the distinct pairs obtained from the final two terms of occurrences of patterns $\tau_1 \cdots \tau_{m-2}m(m-1)$ and $\tau_1 \cdots \tau_{m-2}(m-1)m$ in a permutation, respectively. By a simple involution in terms of permutation diagrams we will prove their equidistribution over the symmetric group. As special case we derive a one-to-one correspondence between permutations which avoid each of the patterns $\tau_1 \cdots \tau_{m-2}m(m-1) \in \mathcal{S}_m$ and such ones which avoid each of the patterns $\tau_1 \cdots \tau_{m-2}(m-1)m \in \mathcal{S}_m$. For $m = 3$ this correspondence coincides with the bijection given by Simion and Schmidt in their famous paper on restricted permutations.

1 INTRODUCTION

In recent time much work has been done investigating permutations with restrictions on the patterns they contain. Given a permutation $\pi \in \mathcal{S}_n$ and a permutation $\tau \in \mathcal{S}_m$, an *occurrence of τ in π* is an integer sequence $1 \leq i_1 < i_2 < \dots < i_m \leq n$ such that the letters of the word $\pi_{i_1}\pi_{i_2} \cdots \pi_{i_m}$ are in the same relative order as the letters of τ . In this context, τ is called a *pattern*. If there is no occurrence we say that π *avoids* τ , or alternatively, π is τ -*avoiding*. We write $\mathcal{S}_n(\tau)$ to denote the set of τ -avoiding permutations in $\mathcal{S}_n$, and more general, $\mathcal{S}_n(T)$ for the set of all permutations of length n which avoid each pattern of T .

A fundamental problem concerning pattern-avoiding permutations is that of discovering explicit bijections between the sets $\mathcal{S}_n(T)$ and $\mathcal{S}_n(T')$ provided they have the same number of elements. The first correspondence of that kind was presented by Simion and Schmidt [10]. It proves the well-known fact that $|\mathcal{S}_n(123)| = |\mathcal{S}_n(132)|$. We will generalize this result.

In [8] and [9], we have used the diagram of a permutation to study certain forbidden patterns. For a permutation $\pi \in \mathcal{S}_n$, we obtain its *diagram* from the $n \times n$ array representation of π by shading, for each dot, the cell containing it and the squares that are due south and due east

Date: February 28, 2003

of it. Each square left unshaded we call a *diagram square*. By the construction, the connected components of all diagram squares form Young diagrams. For a diagram square, its *rank* is defined to be the number of dots northwest of it. Clearly, connected diagram squares have the same rank.

In this paper, we use permutation diagrams again. Section 2 demonstrates that the last two terms of occurrences of patterns $\tau \in \mathcal{S}_m$ with $\tau_{m-1}\tau_m = m(m-1)$ and $\tau_{m-1}\tau_m = (m-1)m$, respectively, are closely related to the diagram squares. It is an essential property of the permutation statistics which count the number of these pairs that they have the same distribution over the symmetric group. To prove this fact, a bijection is established in Section 3 which respects these statistics. In particular, it will be shown that there are as many permutations in $\mathcal{S}_n$ which avoid each pattern $\tau \in \mathcal{S}_m$ with $\tau_{m-1} = m$ and $\tau_m = m-1$ as permutations which avoid each pattern $\tau \in \mathcal{S}_m$ with $\tau_{m-1} = m-1$ and $\tau_m = m$. For $m = 3$ the correspondence coincides with Simion-Schmidt's bijection. We conclude with some final remarks done in Section 4.

2 DIAGRAMS AND OCCURRENCES OF PATTERNS

For $m \geq 2$ define the pattern sets

$$A_m = \{\tau \in \mathcal{S}_m : \tau_{m-1} = m, \tau_m = m-1\} \quad \text{and} \quad B_m = \{\tau \in \mathcal{S}_m : \tau_{m-1} = m-1, \tau_m = m\}.$$

For a permutation $\pi \in \mathcal{S}_n$, denote by $\mathbf{a}_m(\pi)$ resp. $\mathbf{b}_m(\pi)$ the number of distinct pairs (i, j) with $1 \leq i < j \leq n$ such that i, j are the final two terms of an occurrence of a pattern belonging to A_m and B_m , respectively, in π . If $\mathbf{a}_m(\pi) = 0$ then π avoids each pattern of A_m ; analogously, $\mathbf{b}_m(\pi) = 0$ means the avoidance of every pattern of B_m . Clearly, $\mathbf{a}_2(\pi)$ counts the inversions in π while $\mathbf{b}_2(\pi)$ says how often π contains the pattern 12. (Apart from $m = 2$, the numbers $\mathbf{a}_m(\pi)$ and $\mathbf{b}_m(\pi)$, respectively, do not coincide in general with the total numbers of occurrences of patterns contained in A_m or B_m .)

For example, the occurrences of 1243 in $\pi = 7\ 1\ 4\ 2\ 6\ 3\ 5 \in \mathcal{S}_7$ are the sequences $(2, 3, 5, 7)$, $(2, 4, 5, 6)$, and $(2, 4, 5, 7)$; $(3, 4, 5, 7)$ is the only occurrence of 2143. Furthermore, π contains once the pattern 1234 and avoids 2134. Hence $\mathbf{a}_4(\pi) = 2$ and $\mathbf{b}_4(\pi) = 1$.

The number $\mathbf{a}_m(\pi)$ can immediately be read off from the ranked diagram of π .

Proposition 1 *Let $\pi \in \mathcal{S}_n$ be a permutation. Then $\mathbf{a}_m(\pi)$ equals the number of diagram squares of rank at least $m-2$. In particular, π avoids each pattern of A_m if and only if every diagram square is of rank at most $m-3$.*

Proof. It follows from the construction that any diagram square (i, j) of rank at least $m-2$

corresponds to an occurrence of a pattern of A_m whose final terms are just i, k where $\pi_k = j$. $\square$

By definition, the number $\mathbf{b}_m(\pi)$ counts the number of non-inversions on the letters of π which are greater than at least $m - 2$ letters to their left. (Here a pair (i, j) is called a *non-inversion* if $i < j$ and $\pi_i < \pi_j$.) It is easy to see that all informations about a permutation are encoded in the diagram squares of rank at most $m - 3$ and the pairs just counted by $\mathbf{b}_m(\pi)$.

Proposition 2 *Any permutation $\pi \in \mathcal{S}_n$ can completely be recovered from the diagram squares having rank at most $m - 3$ and the pairs (i, j) arising from the final two terms of an occurrence of a pattern belonging to B_m .*

Proof. By the diagram construction, to know all the diagram squares of rank at most $m - 3$ means to know the positions of all elements π_i for which there are at most $m - 3$ integers $j < i$ with $\pi_j < \pi_i$. As mentioned before, the pairs (i, j) obtained from the end of any occurrence of a pattern of B_m are exactly the non-inversions on the set of integers k for which π_k exceeds at least $m - 2$ elements on its left. Thereby the positions of these elements are also uniquely determined. $\square$

The proof implies the following procedure for constructing a permutation from its diagram squares of rank at most $m - 3$ and the final terms of occurrences of patterns belonging to B_m in π .

For some integer $m \geq 2$, let D be a set of squares and O a set of integer pairs that are obtained from a permutation $\pi \in \mathcal{S}_n$ as diagram squares of rank at most $m - 3$ and occurrence terms as described, respectively. First represent the elements of D as white squares in an $n \times n$ array, shaded otherwise. Row by row, put a dot in the leftmost shaded square such that there is exactly one dot in each column. Each dot having more than $m - 3$ dots northwest is being deleted. Then arrange the missing dots such that O is precisely the set of non-inversions on these dots, that is, the dot contained in the i th row lies strictly to the left of the dot contained in the j th row if and only if $(i, j) \in O$.

An efficient way to arrange the dots is the following one. Let $r_1 < r_2 < \dots < r_s$ be the indices of rows containing no dot, and $c_1 > c_2 > \dots > c_s$ the indices of columns without a dot. Furthermore, let e_i be the number of pairs in O whose first component equals r_i . For $i = 1, \dots, s$, set $c'_i = c_{e_i+1}$, delete c_{e_i+1} from the sequence c , and renumber the sequence terms. Put the missing dots in the squares (r_i, c'_i) where $1 \leq i \leq s$. Note that the second component of the pairs contained in O is of no importance for this procedure. It suffices to know the first components and the number of their appearance in O .

Example 3 Let $\pi = 38510241967 \in \mathcal{S}_{10}$ and $m = 5$. The left-hand array shows the ranked

permutation diagram of π . All the occurrences of patterns of B_5 end with $(9, 10)$. Thus we obtain:




Figure 1 Recovering of a permutation

Red dots represent the elements of π which exceed at most two elements on their left. Note that $(9, 10)$ is the only non-inversion on the set of elements represented by black dots in the right-hand array. (The sorting routine yields $c' = (10, 9, 6, 7)$ since $e = (0, 0, 1, 0)$.)

3 THE BIJECTION

The properties of permutation diagrams given in the previous section are essential for the construction of a bijection Φ_m whose aim it will be to prove

Theorem 4 *We have $|\{\pi \in \mathcal{S}_n : \mathbf{a}_m(\pi) = k\}| = |\{\pi \in \mathcal{S}_n : \mathbf{b}_m(\pi) = k\}|$ for all n and k .*

Let $\pi \in \mathcal{S}_n$ be a permutation, and let D be the set of its ranked diagram squares. Define $\sigma = \Phi_m(\pi)$ to be the permutation whose diagram squares of rank at most $m - 3$ coincide with the elements of D of rank at most $m - 3$, and for which the first components of the pairs obtained as final terms of an occurrence of a pattern of B_m are just the row indices of the squares in D having rank at least $m - 2$.

Before we will analyse this map, we give an example.

Example 5 Consider $\pi = 3\ 8\ 5\ 10\ 2\ 4\ 1\ 9\ 6\ 7 \in \mathcal{S}_{10}$ again. For $m = 5$, the map Φ_m takes π to the permutation $\sigma = 3\ 8\ 5\ 9\ 2\ 4\ 1\ 6\ 10\ 7$:




Figure 2 Bijection Φ_5 , applied to $\pi = 3\ 8\ 5\ 10\ 2\ 4\ 1\ 9\ 6\ 7$

The diagram squares having rank at most 2 coincide for π and σ . From the row indices of the diagram squares of π whose rank is at least 3 we obtain the first component of each pair

arising from the final terms of all occurrences of patterns of B_5 in σ . Exactly, we have $O = \{(4, *), (8, *), (8, *)\}$. The construction of σ 's complete array is done as described following Proposition 2. With the notations introduced there, we have $r = (4, 8, 9, 10)$, $c = (10, 9, 7, 6)$, $e = (1, 2, 0, 0)$, and hence $c' = (9, 6, 10, 7)$.

As discussed above, the equality of the diagram squares having rank at most $m - 3$ for π and $\sigma = \Phi_m(\pi)$ means that $\sigma_i = \pi_i$ for every i for which there are at most $m - 3$ integers $j < i$ with $\pi_j < \pi_i$. In particular, π and σ coincide in the $m - 2$ first letters. By diagram construction, each white square of rank greater than $m - 3$ is just a pair (i, π_j) for which there are at least $m - 2$ integers $k < i$ with $\pi_k < \pi_j$. Obviously, we have $i < j$ and $\pi_j < \pi_i$. Hence both π_i and π_j are elements exceeding at least $m - 2$ elements on their left. Consequently, the map Φ_m is well-defined, and bijective by Proposition 2 and the remarks done following it.

It is easy to see that Φ_m yields the equidistribution of $\mathbf{a}_m$ and $\mathbf{b}_m$ over the symmetric group.

Proposition 6 *Let $\pi \in \mathcal{S}_n$ and $\sigma = \Phi_m(\pi)$, for any $m \geq 2$. Then $\mathbf{a}_m(\pi) = \mathbf{b}_m(\sigma)$.*

Proof. By Proposition 1, every pair (i, j) which arises from the two final terms of an occurrence of a pattern of A_m in π corresponds to a diagram square of π having rank at least $m - 2$, namely (i, π_j) . It follows immediately from the definition of Φ_m that there is an occurrence of a pattern of B_m in σ which ends with (i, k) where k depends on j . $\square$

Remarks 7

- a) By the proof, every occurrence of a pattern of A_m in π corresponds in a one-to-one fashion to an occurrence of a pattern belonging to B_m in $\Phi_m(\pi)$ where both sequences coincide in the $(m - 1)$ st term. Consequently, Φ_m is even an involution, and we have $\mathbf{b}_m(\pi) = \mathbf{a}_m(\Phi_m(\pi))$ for all $\pi \in \mathcal{S}_n$.
- b) The bijection Φ_m has the advantage of fixing precisely the intersection of the sets $\mathcal{S}_n(A_m)$ and $\mathcal{S}_n(B_m)$.
- c) The map Φ_2 simply takes a permutation $\pi \in \mathcal{S}_n$ to $\sigma \in \mathcal{S}_n$ with $\sigma_i = n + 1 - \pi_i$. Note that we have to arrange all n dots by the procedure given above. Here e_i equals the number of diagram squares in the i th row, or, equivalently, the number of integers j satisfying $i < j$ and $\pi_i > \pi_j$. The sorting routine yields the permutation $\sigma = c'$ whose occurrences of pattern 12 are just the inversions of π .

The case $k = 0$ of Theorem 4 means the *Wilf equivalence* of the pattern sets A_m and B_m , that is, there are as many permutations in $\mathcal{S}_n$ which avoid each pattern of A_m as those which avoid each pattern of B_m . An analytical proof of this result was given in [6].

Corollary 8 For each $m \geq 2$, the sets A_m and B_m are Wilf equivalent.

For $\pi \in \mathcal{S}_n(A_m)$ the construction of $\sigma = \Phi_m(\pi)$ is even more simple. By Proposition 1, every diagram square of π is of rank at most $m - 3$. Therefore the bijection works as follows. Set $\sigma_i = \pi_i$ if there are at most $m - 3$ integers $j < i$ satisfying $\pi_j < \pi_i$. Then arrange the remaining elements in decreasing order.

For example, the permutation $\pi = 2\ 6\ \underline{7}\ 1\ \underline{3}\ \underline{4}\ \underline{5} \in \mathcal{S}_7$ avoids both 1243 and 2143. We obtain $\Phi_4(\pi) = 2\ 6\ \underline{7}\ 1\ \underline{5}\ \underline{4}\ \underline{3} \in \mathcal{S}_7(B_4)$. (All elements which exceed at least two elements on their left are underlined.)

Particularly for $m = 3$, all the left-to-right minima of π are preserved. (A *left-to-right minimum* of a permutation π is an element π_i which is smaller than all elements to its left, i.e., $\pi_i < \pi_j$ for every $j < i$.) The other elements of π are being put on the free positions, in decreasing order. This is precisely the description of the bijection between $\mathcal{S}_n(132)$ and $\mathcal{S}_n(123)$ proposed by Simion and Schmidt in [10, Prop. 19].

4 AN OPEN QUESTION

The number of elements of $\mathcal{S}_n(B_m)$ (or, equivalently, $\mathcal{S}_n(A_m)$) was determined in [1]. Similar to the studies of the number of occurrences of certain patterns which are intensified in last time, we may ask for the number $|\{\pi \in \mathcal{S}_n : \mathbf{a}_m(\pi) = k\}|$ for any positive integer k . (By Theorem 4, the problem of determining $|\{\pi \in \mathcal{S}_n : \mathbf{b}_m(\pi) = k\}|$ is equivalent.) As shown in Proposition 1, this number just counts the permutations in $\mathcal{S}_n$ whose diagram has exactly k squares of rank at least $m - 2$. Here we only consider the special case $m = 3$ and $k = 1$.

Our proof uses tunnels in Dyck paths which were introduced very recently by Elizalde [4]. Recall that a *Dyck path* of length $2n$ is a lattice path in $\mathbb{Z}^2$ between $(0, 0)$ and $(2n, 0)$ consisting of up-steps $[1, 1]$ and down-steps $[1, -1]$ that never falls below the x -axis. For any Dyck path d , a *tunnel* is defined to be a horizontal segment between two lattice points of d that intersects d only in these two points, and stays always below d . The *length* and *height* of a tunnel are measured in the lattice. Figure 3 shows a tunnel (drawn with a red line) of length 4 and height 2.




Figure 3 A tunnel in a Dyck path

In all Dyck paths of length $2n$ there are $\binom{2n-1}{n-3}$ tunnels of positive height and length at least 4. To verify this, note that there are nC_n tunnels in all since every tunnel is associated with an up-step, and the number of Dyck paths of length $2n$ equals the n th *Catalan number* $C_n = \frac{1}{n+1} \binom{2n}{n}$.

The tunnels of height zero correspond precisely to *returns*, that is, down-steps landing on the x -axis. By [3], the total number of returns in Dyck paths of length $2n$ is equal to $\frac{3}{2n+1} \binom{2n+1}{n-1}$. Each tunnel of length 2 and positive height is just the connection line of a *high peak*. (A *high peak* of a Dyck path is an up-step followed by a down-step whose common lattice point is at a level greater than 1.) Their number was also given in [3]; it equals $\binom{2n-1}{n-2}$.

Proposition 9 *We have $|\{\pi \in \mathcal{S}_n : \mathbf{a}_3(\pi) = 1\}| = \binom{2n-1}{n-3}$ for all n .*

Proof. We have to count the permutation diagrams having exactly one square of rank $r \geq 1$. Every such diagram arises from a permutation diagram having only squares of rank zero (note that the permutations whose diagram satisfies this condition are just the 132-avoiding ones) by adding the additional white square (i, j) such that $i + j \leq n + r$, the number of shaded squares (i, k) with $1 \leq k < j$ is equal to r , and the number of shaded squares (k, j) with $1 \leq k < i$ is equal to r , too. In addition, the $r \times r$ subarray consisting of the squares (i', j') with $i - r \leq i' \leq i - 1$ and $j - r \leq j' \leq j - 1$ must be the diagram of any permutation in $\mathcal{S}_r(132)$. This condition is equivalent to $i' + j' + r + 2 \leq i + j$ for all diagram squares (i', j') contained in the subarray. Consider now the Dyck path that goes from the lower-left corner to the upper-right corner of the array, and travels along the boundary of the connected component of all diagram squares of rank zero.




Figure 4 Correspondence between diagram square of rank $r \geq 1$ and Dyck path tunnel

Any square (i, j) satisfies the above conditions if and only if the line that connects the path step contained in the i th row (up-step) with the path step contained in the j th column (down-step) is a tunnel of length $2r + 2 \geq 4$ and height at least 1. $\square$

Remark 10 Thomas [11] gives the following alternative combinatorial proof of Proposition 9 using the permutation statistic $\mathbf{b}_3$:

Let $\pi \in \mathcal{S}_n$ satisfy $\mathbf{b}_3(\pi) = 1$. Furthermore, let (i, j) be the final terms of any occurrence of the pattern 123 in π . Consider now the permutation $\sigma \in \mathcal{S}_n$ which arises from π by exchanging π_i with π_j . It is easy to see that σ avoids 123. What can we say about the elements σ_i and σ_j ? They are successive right-to-left maxima of σ , and there is at least one element to the left of σ_i which is smaller than σ_j . (An element is called a *right-to-left maximum* of a permutation if it exceeds all the elements on its right.) In fact, for any $\sigma \in \mathcal{S}_n(123)$ these two properties characterize the pairs (σ_i, σ_j) whose transposition yields a permutation π for which $\mathbf{b}_3(\pi) = 1$.

Consequently, we want to count right-to-left maxima of 123-avoiding permutations for which the set of elements to their right is not a complete interval $[1, k]$ for some k or the empty set. In [5], Krattenthaler describes a bijection between 123-avoiding permutations in $\mathcal{S}_n$ and Dyck paths of length $2n$ having the property that any right-to-left maximum of the kind we consider corresponds to a valley at a level greater than zero. (A *valley* of a Dyck path is a down-step followed by an up-step.) By [3], these valleys are just counted by $\binom{2n-1}{n-3}$.

For comparison, Noonan [7] proved that the number of permutations in $\mathcal{S}_n$ containing 123 exactly once is given by $\frac{3}{n} \binom{2n}{n-3}$ while Bóna [2] showed that there are $\binom{2n-3}{n-3}$ permutations in $\mathcal{S}_n$ having exactly one 132-subsequence. By [8, Th. 5.1], the latter permutations are characterized to be such ones having exactly one diagram square of rank 1 and only rank 0 squares otherwise.

REFERENCES

1. E. Barcucci, A. Del Lungo, E. Pergola, and R. Pinzani, Permutations avoiding an increasing number of length-increasing forbidden subsequences, *Discrete Math. Theor. Comput. Science* **4** (2000), 31-44.
2. M. Bóna, Permutations with one or two 132-subsequences, *Discrete Math.* **181** (1998), 267-274.
3. E. Deutsch, Dyck path enumeration, *Discrete Math.* **204** (1999), 167-202.
4. S. Elizalde, Fixed points and excedances in restricted permutations, preprint, math.CO/0212221.
5. C. Krattenthaler, Permutations with restricted patterns and Dyck paths, *Adv. Appl. Math.* **27** (2001) 510-530.
6. T. Mansour, Permutations with forbidden patterns, Ph.D. Thesis, University of Haifa, 2001.
7. J. Noonan, The number of permutations containing exactly one increasing subsequence of length three, *Discrete Math.* **152** (1996), 307-313.
8. A. Reifegerste, On the diagram of 132-avoiding permutations, preprint, math.CO/0208006.
9. A. Reifegerste, On the diagram of Schröder permutations, *Electr. J. Combin.* **9(2)** (2003), R8.
10. R. Simion and F. W. Schmidt, Restricted Permutations, *Europ. J. Combin.* **6** (1985), 383-406.
11. H. Thomas, personal communication.